





How to prepare for your
Colonoscopy
with HalfLytely and Bisacodyl
Tablets

Follow only these instructions (disregard the instructions that come with the kit). Please read over these instructions carefully as soon as you receive them.

Remember, your appointment could be rescheduled or cancelled if:

- You do not finish your prep or appropriately follow the prep instructions below.
- You consume any food or drink within 3 hours of your arrival time.
- You do not arrive for your appointment on time.
- You do not have a ride home. Medications will be given to you at the time of your colonoscopy that will impair your judgment and reflexes. You will not be able to drive yourself, and you will only be able to use a taxi or bus service if you have a responsible adult age 18 year or older to accompany you.
- You have a change in your health insurance coverage. Your new plan may require us to obtain authorization before performing your colonoscopy, a process that can take some time. Please notify us of any changes to your insurance immediately.

## One week before your colonoscopy...

- 1. **Fill your prescription for HalfLytely and Bisacodyl Tablets if you have not already.** If your pharmacy does not contact you regarding this prescription, please call them to verify if/when it is ready of pickup. If no prescription was received, is not covered by your insurer, or is too costly, please call (203) 281-4463, ext. 5152 to speak with our prescription line.
- 2. Purchase items for the clear liquid diet:
  - o Clear broth or bouillon
  - Soda, Gatorade (yellow or clear) or clear fruit juice (apple, white grape, lemonade)
  - o Popsicles (no purple, red, or orange) or italian ice (lemon or lime)
  - Jell-O (no purple, red, or orange)
  - o Coffee or Tea (no milk, cream, or dairy).
- 3. **Arrange for transportation to and from your colonoscopy.** You will not be allowed to drive yourself. You may not use a taxi or bus service unless you have a responsible adult 18 years or older to accompany you.
- 4. Please refer to the instructions provided by the office regarding preprocedure medication changes. If you have any questions, please contact the office.
- 5. Stop taking iron supplements (multivitamins containing iron are OK).
- 6. Must purchase a bottle of Infant's Mylicon (Only if box is checked)

### One day before your colonoscopy...

- 1. **Upon waking up**, begin the clear liquid diet:
  - o No solid food until after your procedure
  - Water
  - Clear broth or bouillon
  - o Soda, Gatorade (yellow or clear) or clear fruit juice (apple, white grape, lemonade)
  - o Popsicles (no purple, red, or orange) or italian ice (lemon or lime)
  - Jell-O (no purple, red, or orange)
  - o Coffee or Tea (no milk, cream, or dairy).
  - o There is no limit to the amount of clear liquids you drink during the day.

# 2. In the morning

o Fill the HalfLytely container with water and refrigerate

#### At noon

o Take 1 Bisacodyl tablet with a sip of water

# 4. At 3pm in the afternoon

- Drink half of the HalfLytely solution. Consume 8 ounces every 10-15 minutes.
- o Refrigerate the remaining half for the morning.

## On the day of your colonoscopy...

1.	No solid food until after your procedure
2.	4 hours prior to your arrival time:
	<ul> <li>Stir up the remaining HalfLytely solution. Consume it within 15-30 minutes.</li> </ul>
	<ul> <li>Add 6ml Infant's Mylicon to the 8oz of water and consume (Only if box is checked)</li> </ul>
3.	3 hours prior to your arrival time stop consuming anything by mouth (liquids or solids).
4.	Make sure to arrive on time for your appointment.