



PACT Gastroenterology Center
Hamden • Milford • Guilford



How to prepare for your Colonoscopy with Sutab Kit

Follow only these instructions (disregard the instructions that come with the kit). Please read over these instructions carefully as soon as you receive them.

Remember, your appointment could be rescheduled or cancelled if:

- **You do not finish your prep or appropriately follow the prep instructions below.**
- **You consume any food or drink within 3 hours of your arrival time.**
- **You do not arrive for your appointment on time.**
- **You do not have a ride home.** Medications will be given to you at the time of your colonoscopy that will impair your judgment and reflexes. You will not be able to drive yourself, and you will only be able to use a taxi or bus service if you have a responsible adult age 18 year or older to accompany you.
- **You have a change in your health insurance coverage.** Your new plan may require us to obtain authorization before performing your colonoscopy, a process that can take some time. Please notify us of any changes to your insurance immediately.

One week before your colonoscopy...

1. **Fill your prescription for Sutab if you have not already.** If your pharmacy does not contact you regarding this prescription, please call them to verify if/when it is ready of pickup. If no prescription was received, is not covered by your insurer, or is too costly, please call (203) 281-4463, ext. 5152 to speak with our prescription line.
2. **Purchase items for the clear liquid diet:**
 - Clear broth or bouillon
 - Soda, Gatorade (yellow or clear) or clear fruit juice (apple, white grape, lemonade)
 - Popsicles (no purple, red, or orange) or italian ice (lemon or lime)
 - Jell-O (no purple, red, or orange)
 - Coffee or Tea (no milk, cream, or dairy).
3. **Arrange for transportation to and from your colonoscopy.** You will not be allowed to drive yourself. You may not use a taxi or bus service unless you have a responsible adult 18 years or older to accompany you.
4. **Please refer to the instructions provided by the office regarding preprocedure medication changes. If you have any questions, please contact the office.**
5. Stop taking iron supplements (multivitamins containing iron are OK).
6. **Must purchase a bottle of Infant's Mylicon (Only if box is checked)**

Tel: 203-281-4463

www.pactmd.com

Fax: 203-287-2930

2200 Whitney Avenue Suites 330 & 360
Hamden CT 06518

40 Commerce Park
Milford CT 06460

1591 Boston Post Road Suite 206
Guilford CT 06437

One day before your colonoscopy...

1. **Upon waking up**, begin the clear liquid diet:
 - No solid food until after your procedure
 - Water
 - Clear broth or bouillon
 - Soda, Gatorade (yellow or clear) or clear fruit juice (apple, white grape, lemonade)
 - Popsicles (no purple, red, or orange) or italian ice (lemon or lime)
 - Jell-O (no purple, red, or orange)
 - Coffee or Tea (no milk, cream, or dairy).
 - There is no limit to the amount of clear liquids you drink during the day.

2. **At 3pm in the afternoon**
 - Open the bottle of 12 Sutab tablets.
 - Fill the provided container up to the fill line with 16 ounces of water. Swallow each of the 12 tablets with a sip of water every 2-3 minutes. Consume all 12 pills and 16 ounces of water over 30 minutes. Please feel free to drink additional water as needed.
 - Approximately 60 minutes after the last tablet is ingested, fill the provided container a second time with 16 ounces of water. Drink the entire amount over 30 minutes.
 - Approximately 30 minutes after finishing the second container of water, fill the provided container a third time with 16 ounces of water. Drink the entire amount over 30 minutes.
 - Add 6ml Infant's Mylicon to the last 16oz of water (Only if box is checked)

On the day of your colonoscopy...

1. **No solid food until after your procedure**
2. **6 hours prior to your arrival time:**
 - Open the bottle of 12 Sutab tablets.
 - Fill the provided container up to the fill line with 16 ounces of water. Swallow each of the 12 tablets with a sip of water every 2-3 minutes. Consume all 12 pills and 16 ounces of water over 30 minutes. Please feel free to drink additional water as needed.
 - Approximately 60 minutes after the last tablet is ingested, fill the provided container a second time with 16 ounces of water. Drink the entire amount over 30 minutes.
 - Approximately 30 minutes after finishing the second container of water, fill the provided container a third time with 16 ounces of water. Drink the entire amount over 30 minutes.
 - Add 6ml Infant's Mylicon to the last 16oz of water (Only if box is checked)
3. **3 hours prior to your arrival time stop consuming anything by mouth (liquids or solids).**
4. Make sure to arrive on time for your appointment.